



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

Healing Body & Mind

Once you have the procedure you waited a long time for, it is time to be patient and wait, again, for your body to heal. Unfortunately, a surgeon's scalpel is not a magic wand and healing will occur at different rates for different people. You must allow yourself to heal and be aware it is natural to be impatient for the results and a little anxious because you often look worse before you look better. Being aware this can happen will help you understand these normal feelings.

Sharing your experience with friends and family can help give you a support system during your recovery, but realize they may unintentionally make you question your decision with concerns during your recovery. While you may think that you are looking pretty rough around the edges, we may tell you that you are healing beautifully. Trust us. We will share with you if there is a concern, so if we tell you that everything is healing normally, it is.

Your healing will depend on many things such as your general health, your willingness to follow instructions, and your mental attitude toward recovery. While I can perform the surgery, I cannot "heal" you. It is up to you to be an active participant in your recovery process to help your body heal the best that it can. Following all of our instruction is very important, as is working with us to address any complications that may arise. Even surgeries that are done exactly right, can have complications during recovery. It has to do with the human factor. We are not machines and every person can react differently and heal differently even when the surgery is done exactly the same. Every surgeon has unexpected results from time to time.

It is important you approach your surgery and especially your recovery with the mindset that we are a team, and you are an integral part of that team. We must trust each other to be working for a common goal, your successful result. As the surgeon, I enjoy my work and strive to achieve an ideal result during every surgery for every patient. I have rigorous standards that my staff must meet in order to be involved in your care. I am looking forward to working with you to achieve a great result and do not anticipate any post-operative problems. I will do everything I can to make sure that you are happy with your result.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

Your Anesthesia Experience

Anytime you undergo anesthesia, your general health condition must be considered. Depending on your age and the results of your history and physical, you may need pre-operative testing such as blood work, chest x-ray, or EKG. This is done to ensure that your current health does not put you at any additional unnecessary risk while under anesthesia.

There Are Four Levels Of Sedation That We Offer:

Local Sedation

A local anesthetic agent such as Novocaine, Xylocaine, or Marcaine is introduced in to the tissue through injection and produces a numbness that allows surgery to be carried out with little to no discomfort while the patient is awake.

Oral Sedation

Local sedation is used in conjunction with Valium (sedative), Percocet (pain medication), and Phenergan (anti-nausea medication) which are ingested orally. You will experience little to no discomfort; be aware and conscious but fairly relaxed; and responsive to voice.

Conscious Sedation with Registered Nurse and the physician

Versed or Fentanyl is administered through an intravenous line by our nurse. This level is a little deeper state of sedation. You will be very sleepy but still responsive to more forceful amounts of voice and touch. Some refer to this type of sedation as Twilight.

Conscious Sedation with a Certified Registered Nurse Anesthetist

The last type of anesthesia offered is administered through an intravenous line given by a Certified Registered Nurse Anesthetist. This level can allow you to go into a deeper state of conscious sedation where you require more stimulation through touch and voice to become responsive. With this level of sedation, you will be much less aware of what is going on during the procedure than the other levels.

General Anesthesia with a Certified Registered Nurse Anesthetist or Anesthesiologist

This type of anesthesia offered is done through both an Intravenous line and an endotracheal tube (breathing tube down the windpipe). This level will take you to deepest level of sedation. You will not be aware of anything and will be in a deep sleep. We use the same medication as 3 and 4 options and also some anesthesia gases.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

We ask that you strictly avoid food and liquids 6 hours before your procedure. This is done for your protection to empty your stomach and prevent aspiration of food or fluid from the stomach into the lungs during anesthesia. Importantly, prior to this point we would like you to drink 64 ounces of fluid with solutes in it like Gatorade, Grape Juice, v8 etc. (Gatorade is probably the best). Your urine should be clear by the time you go to bed. If you are just having local anesthesia with your procedure and no sedation with the physician, you may be allowed to have something light before your procedure BUT YOU SHOULD ASK THE PHYSICIAN BEFORE DOING THIS.

Either prior to or on the day of surgery, your anesthesiologist or nurse will have questions for you regarding your health, height, weight and past anesthesia experiences. Complete and honest answers are required to assist your anesthesiologist or nurse in planning and administering the safest level of anesthetic possible. As you wake up from your anesthesia, your vitals will still be monitored closely for a half an hour. Often you will be given additional medications for discomfort and/or nausea. Once you are awake and aware, you may be released to a responsible adult to go home with you. Please follow the post-operative instructions closely regarding supervision requirements and activity restrictions. Drinking lots of fluids will help rid your body of the medications used for anesthesia more quickly.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

Pre-Procedure Instructions – General

The following instructions should be followed closely except when overruled by specific procedural instructions.

Two Weeks Prior to Your Procedure and After Procedure

NO ASPIRIN or medicines that contain aspirin* since it interferes with normal blood clotting.

NO IBUPROFEN or medicines contain ibuprofen* as it interferes with blood clotting.

Please DISCONTINUE ALL HERBAL MEDICATIONS* as many have side effects that could complicate a surgical procedure by inhibiting blood clotting, affecting blood pressure, or interfering with anesthetics.

Please DISCONTINUE ALL DIET PILLS whether prescription, over-the-counter or herbal as many will interfere with anesthesia and can cause cardiovascular concerns.

NO “MEGADOSES” OF VITAMIN E (which is anything greater than 40IU), but a multiple vitamin that contains E is just fine.

NO SMOKING because nicotine reduces blood flow to the skin and can cause significant complications during healing at least 2 weeks before and after your procedure.

You may take Tylenol or generic forms of this drug. These do not interfere with blood clotting or healing.

Start taking a multivitamin each day and continue taking through your recovery. The healthier you are, the quicker your recovery will be.

One Week Prior to Your Procedure

DO NOT take or drink any alcohol or drugs for one week prior to your procedure and one week after your procedure as these can interfere with anesthesia and affect blood clotting.

DO NOT color your hair from this point on and for one month after your procedure.

If your skin tolerates, use a germ-inhibiting soap for bathing, such as Dial, Safeguard, or Lever 2000 for at least the week before your procedure.

DO report any signs of cold, infection, boils, or pustules appearing before your procedure.

DO NOT take any cough or cold medications without permission.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

DO arrange for a responsible adult to drive you to and from the facility on the day of your procedure, since you will not be allowed to leave on your own. We will have a caretaker form for your caretaker to sign.

DO arrange for a responsible individual to spend the first 24 hours with you, since you CANNOT be left alone.

Day Before Your Procedure and Morning of Your Procedure

We may prescribe Clonidine to you that you should take the day before your procedure; ½ to 1 tablet in the morning and again at night; and that will continue for 48 hours after your procedure.

If you are having oral or conscious sedation you should have nothing to eat or drink (but you can have a little sip of water to take your usual medications) 6 hours before your procedure. Also, no gum, candy, mints or coffee the morning of your procedure. Do not sneak anything as this may endanger you. If you are just having your procedure under local anesthesia in our office you can eat something light up until your procedure.

If you are on regular medications, please clear these with Your physician. Otherwise, please make sure to bring all of the medications prescribed by Your physician with you on the day of the procedure and your own medications that you regularly take.

DO take a thorough shower with your germ-inhibiting soap the night before and the morning of your procedure. Shampoo your hair the morning of your procedure. This is to decrease the bacteria on the skin and thereby decrease the risk of infection.

DO NOT apply any of the following to your skin, hair or face the morning of your procedure: make-up, creams, lotions, hair gels, sprays, perfumes, powder, or deodorant. Using any of these products will add bacteria to the skin and increase the risk of infection.

You may brush your teeth the morning of your procedure but do not drink anything unless you are having the procedure in our office under local anesthesia without oral or conscious sedation.

DO NOT wear contacts to your procedure. If you do wear glasses, bring your eyeglass case.

DO wear comfortable, loose-fitting clothes that do not have to be put on over your head. The best thing to wear home is a button-up top and pull on pants. You will want easy-to-slip-on flat shoes.

DO NOT bring any valuables or wear any jewelry (no rings, earrings, chains, toe rings, other metal piercings or watches). We will need to tape wedding rings if worn.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

You must have an adult drive for you – to and from your procedure. Please note that a cab or bus driver will not be allowed to take you home after your procedure. On arrival, be sure we know your driver's name, phone numbers, and how we will be able to reach them.

If you are not recovering at home, it is very important that we have the number where you will be after your procedure.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

Medications to Avoid

If you are taking any medications on this list, they should be discontinued 2 weeks before and after your procedure and only acetaminophen products, such as Tylenol, should be taken for pain. Most importantly we would like you to avoid high dose vitamin E (anything greater than 40IU), aspirin, anti-inflammatories, herbal medications, supplements (fish oil, omega 3's). All other medications – prescriptions, over-the-counter and herbal medications or supplements– that you are currently taking must be specifically cleared by your Doctor prior to surgery. It is absolutely necessary that all of your current medications be specifically cleared by your Doctor and the nursing staff. **It is okay to eat the foods listed, we just don't want you to eat excessive amounts of the foods listed below.**

Aspirin Medications to Avoid: *Affect blood clotting.*

4-Way Cold Tabs	Aspergum	Cheracol	Equagesic
5-Aminosalicylic Acid	Asprimox products	Choline Magnesium	Etodolac
Acetilsalicylic Acid	Axotal	Trisalicylate	Excedrin products
Actron	Azdone	Choline Salicylate	Fiorgen PF
Adprin-B products	Azulfidine products	Cope	Fiorinal products
Aleve	B-A-C	Coricidin	Flurbiprofen
Alka-Seltzer products	Backache Maximum	Cortisone	Gelpirin
Amigesic Argesic-SA	Strength Relief	Medications	Genprin
Anacin products	Bayer Products	Damason-P	Gensan
Anexsia w/Codeine	BC Powder	Darvon Compound-	Goody's Extra
Arthra-G	Bismatrol products	65	Strength Headache
Arthriten products	Buffered Aspirin	Darvon/ASA	Powders
Arthritis Foundation	Bufferin products	Diclofenac	Halfprin products
products	Buffetts 11	Dipenturn	IBU
Arthritis Pain	Buffex	Disalcid	Indomethacin
Formula	Butal/ASA/Caff	Doan's products	products
Arthritis Strength BC	Butalbital Compound	Dolobid	Isolyl Improved
Powder	Cama Arthritis Pain	Dristan	Kaodene
Arthropan	Reliever	Duragesic	Lanorinal
ASA	Carisoprodol	Easprin	lbuprohm
Asacol	Compound	Ecotrin products	Lodine
Ascriptin products	Cataflam	Empirin products	Lortab ASA



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

Magan	Night-Time	Piroxicam	St. Joseph Aspirin
Magnaprin products	Effervescent Cold	Propoxyphene	Sulfasalazine
Magnesium Salicylate	Norgesic products	Compound products	Supac
Magsal	Norwich products	Robaxisal	Suprax
Marnal	Olsalazine	Rowasa	Synalgos-DC
Marthritic	Orphengesic products	Roxeprin	Talwin
Mefenamic Acid	Orudis products	Saletto products	Triaminicin
Meprobamate	Oxycodonc	Salflex	Tricosal
Mesalamine	Pabalate products	Salicylate products	Trilisate
Methocarbarnol	P-A-C	Salsalate	Tussanil DH
Micrainin	Pain Reliever Tabs	Salsitab	Tussirex products
Mobidin	Panasal	Scot-Tussin Original	Ursinus-Inlay
Mobigesic	Pentasa	5-Action	Vanquish
Momentum	Pepto-Bismol	Sine-off	Wesprin
Mono-Gesic	Percodan products	Sinutab	Willow Bark products
Motrin products	Phenaphen/Codeine	Sodium Salicylate	Zorprin
Naprelan	#3	Sodol Compound	
Naproxen	Pink Bismuth	Soma Compound	

Ibuprofen Medications to Avoid: *Affect blood clotting.*

Acular (ophthalmic)	Haltran	Nabumetone	Rhinocaps
Advil products	Indochron E-R	Nalfon products	Sine-Aid products
Anaprox products	Indocin products	Naprosyn products	Sulindac
Ansaid	Ketoprofen	Naprox X	Suprofen
Clinoril	Ketorolac	Nuprin	Tolectin products
Daypro	lbuprin	Ocufen (ophthalmic)	Tolmetin
Dimetapp Sinus	lbuprofen	Oruvail	Toradol
Dristan Sinus	Meclofenamate	Oxaprozin	Voltaren
Feldene	Meclomen	Ponstel	
Fenoprofen	Menadol	Profenal	
Genpril	Midol-products	Relafen	

Avoid ALL Diet Aids – Including Over-the-Counter & Herbal

Intensify anesthesia, serious cardiovascular effects.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

Tricyclic Antidepressants to Avoid: *Intensify anesthesia, cardiovascular effects.*

Adapin	Clomipramine	Limbitrol products	Sinequan
Amitriptyline	Desipramine	Ludiomil	Surmontil
Amoxapine	Doxepin	Maprotiline	Tofranil
Anafranil	Elavil	Norpramin	Triavil
Asendin	Endep	Nortriptyline	Trimipramine
Aventyl	Etrafon products	Pamelor	Vivactil
	Imipramine	Pertofrane	
	Janimine	Protriptyline	

Other Medication to Avoid: *Affect blood clotting.*

4-Way w/ Codeine	Dicumerol	Miradon	Sparine
A.C.A.	Dipyridamole	Opasal	Stelazine
A-A Compound	Doxycycline	Pan-PAC	Sulfipyrazone
Accutrim	Emagrin	Pentoxyfylline	Tenuate
Actifed	Enoxaparin injection	Persantine	Tenuate Dospan
Anexsia	Flagyl	Phenylpropanolamin e	Thorazine
Anisindione	Fragmin injection	Prednisone	Ticlid
Anturane	Furadantin	Protarnine	Ticlopidine
Arthritis Bufferin	Garlic	Pyrroxate	Trental
BC Tablets	Heparin	Ru-Tuss	Ursinus
Childrens Advil	Hydrocortisone	Salatin	Virbamycin
Clinoril C	Isollyl	Sinex	Vitamin E
Contac	Lovenox injection	Sofarin	Warfarin
Coumadin	Macrochantin	Soltice	
Dalteparin injection	Mellaril		

Salicylate Medications, Foods & Beverages to Avoid: *Affect blood clotting.*

(High concentration of foods to be avoided, you do not need to cut out these foods completely.)

Amigesic (salsalate)	salicylate)	Pamprin (Maximum Pain Relief)	Pabalate
Disalcid (salsalate)	Dolobid (diflunisal)	Mobigesic	Pepto-Bismol (bismuth
Doan's (magnesium	Magsal		



AESTHETIC FACIAL BODY PLASTIC SURGERY
 DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

subsalicylate)	magnesium	Cherries	Pickles
Salflex (salsalate)	salicylate)	Chinese Black Beans	Prunes
Salsalate		Cucumbers	Raspberries
Salsitab (salsalate)	Almonds	Currants	Strawberries
Trilisate (choline	Apples	Garlic	Tomatoes
salicylate +	Apricots	Ginger	Wine
	Blackberries	Grapes	
	Boysenberries		

Vitamins and Herbs to Avoid

Affect blood clotting, affect blood sugar, increase or decrease the strength of anesthesia, rapid heartbeat, high blood pressure, liver damage. Note: Just because it is not of this list does not mean that it is safe to take while preparing for surgery.

Ackee fruit	Devil's club	Goldenseal	Muwort
Alfalfa	Dong Quai root	Gotu Kola	Nem seed oil
Aloe	Echinacea	Grape seed	Onions
Argimony	Ephedra	Guarana	Papaya
Barley	Eucalyptus	Guayusa	Periwinkle
Bilberry	Fenugreek seeds	Hawthorn	Selenium
Bitter melon	Feverfew	Horse Chestnut	St. John's Wort
Burdock root	Fo-ti	Juniper	Valerian/Valerian Root
Carrot oil	Garlic	Kava Kava	"The natural Viagra®"
Cayenne	Ginger	Lavender	Vitamin E
Chamomile	Gingko	Lemon verbena	Willow bark
Chromium	Gingko biloba	Licorice root	Yellow root
Coriander	Ginseng	Ma Huang	Yohimbe
Dandelion root	Gmena	Melatonin	



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

If you are taking anything not on this list, please call the office at (425) 990-3223 to notify us and make sure that it is okay.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

Post-Operative Instructions – General

The following instructions should be followed closely except when overruled by specific procedural instructions. You must follow your surgeon's instructions as indicated for your specific procedure. Notify your physician of any unusual changes in your condition and feel free to call the office with any questions.

You **MUST HAVE AN ADULT DRIVE YOU** home from the facility. You will not be allowed to drive yourself or use public transportation.

After your procedure you **MUST HAVE A RESPONSIBLE ADULT STAY WITH YOU** a minimum of 24 hours. You **CANNOT** be left alone. The 24 hours begin when you are discharged from the office or hospital. Have everything ready at home **PRIOR** to your procedure. Make arrangements for someone to stay with you. Let the person or persons know you cannot be left alone. This is important because of the danger of falling and you may lose the concept of time for the day and overmedicate yourself. Hence, you should always record how much and at what time you take your medications after your procedure.

The effects of anesthesia can persist for 24 hours. You must exercise extreme caution before engaging in any activity that could be harmful to yourself or others.

DRINK fluids to help rid the body of the drugs used in your procedure. If you have straws in the house you will tend to drink more fluids the first few days after your procedure.

Diet may be as tolerated. Eating foods that are bland and soft for the first day or so – foods like after you have had the flu – may be best tolerated. You must eat more than crackers and juice; otherwise you will continue to feel weak and will not heal as well. **REMEMBER** to take the medications with a little something to eat or you will get sick to your stomach.

Please avoid the use of alcoholic beverages for the first 2 weeks after your procedure (it dilates blood vessels and can cause unwanted bleeding) and as long as pain medications are being used (dangerous combination).

Take only medications that have been prescribed by your physician for your postoperative care and take them according to the instructions on the bottle. Your pain medication may make you feel “spacey”; therefore, have someone else give you your medications according to the proper time intervals. You and your caretaker should record how many pills you have, how much and what you are given at each dosage, at what date and time the drugs were given each and every time medications are given to you.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

If you experience any generalized itching, rash, wheezing or tightness in the throat, stop taking all medications and the office immediately if in doubt call 911 , as this may be a sign of a drug allergy.

You can expect moderate discomfort, which should be helped by the pain medications. The greatest discomfort is usually during the first 24 hours. Thereafter, you will find that you require less pain medication.

Call (425) 990-3223 if you have: SEVERE PAIN not responding to pain medication; swelling that is greater on one side than the other; incisions that are RED OR FEVERISH; a FEVER; or if any other questions or problems arise.

Keep any DRESSINGS ON, CLEAN AND DRY. Do not remove them until instructed to do so. There may be some bloody drainage on the dressings. If you have excessive bleeding or the bandages are too tight, call the office immediately.

After your procedure it is important to have a bowel movement within a day or two. If you do not, you may take over the counter laxatives to encourage your bowels to move. If it persists you can call your Doctor anytime or our office.

Minimal activity for the first 48 hours. No house cleaning, furniture rearranging, etc. Relax, be pampered, and let your body heal. The less energy you use on doing things, the more energy your body can focus on healing.

Limit lifting, pulling or pushing for the first 2 weeks. The limit of lifting should be anything under 5 pounds.

Having your head of your bed elevated 45 degrees will help your swelling. This requires a pillow under the small of your back, two pillows under your shoulders and head, and if you have a pillow under each elbow you will relax and stay in position.

You are requested to remain within a reasonable traveling distance of the office for approximately 7-10 days.

Once cleared to shower you may do so every day. We usually allow this 72 hours after your procedure. Please do not use the bathtub for 2 weeks. For the first 2 weeks, you should avoid really hot showers. Lukewarm showers will prevent you from having bleeding, oozing and more swelling.

NO SMOKING for the first 2 weeks before and after your procedure. Any cheating will delay healing.

You may drive once you are off the pain pills and any sedating medications, and when you experience no pain with this activity (you need to be able to react quickly).



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

All surgeries involve some scarring, which can take up to a year to fade. No matter how small they may be, we still want them to heal as well as they are able. Exposing red scars to the sun can cause permanent discoloration. A good sunscreen (SPF 45 or higher) can help and will protect the surrounding tissues that might not feel sunburn developing while the nerves are healing. Sunlight can even reach scars under a swimsuit, so take adequate precautions.

DO NOT use a hot tub for 4 weeks.

AVOID sports or strenuous activities 4 to 6 weeks as your surgeon gives you clearance during your post-operative visits. This is to avoid any unnecessary complications (bleeding, bruising, and swelling).

You may return to work when you feel able and are cleared to do so by your surgeon.

Do not color your hair for 1 month after your procedure.

Feel free to call upon us at any time. We want you to be as comfortable as possible during your healing period.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

Brow Lift Post Procedure Care (Short Form)

Cleaning:

Mixture: 1tsp vinegar to 1 quart of distilled water

1. Clean incisions 3-4 times a day and every 4 hours while asleep.
2. NO crusting, oozing or blood in sutures. **These sutures will be removed 5-7 days post op.** Crusting leads to scarring and infections.
3. Can slow down cleanings as long as you are not crusting up quickly.
4. You should keep Vaseline over treated at all times for the first 7-10 days or until further notified.

Wrapping

1. Wear wrap at all time for 2 weeks, then just at night for 2 months to avoid abnormal swelling.
2. Wear wrap tight for 2 hours, then loose for 1 hour. Keep the wrap at a light compression while sleeping. You do not want to wrap too tightly as this can kill the skin and cause complications.
3. Clean incisions under wrap 3-4 times a day to prevent infection and place Vaseline on the incisions.
4. For areas that are more swollen than others. You can hold pressure on it to flatten it for 5-15 min, 5 times a day. You can use your fingers or use a ace wrap with a piece of tissue the exact same size as the elevation under the ace.

Drain Care:

1. Check the output of each drain separately record and empty each drain every 8 hours. We suggest doing this at 7am, 3pm, and 11pm. You should record how much drainage is in the bulbs with cc or milliliter (ml) increments with the provided measuring cup. The drain is optimally taken out when the output is less than 15-30cc(s) in a 24 hour period. Don't forget to "milk" the drain every few hours to prevent tubing from clotting off. Make sure to let your Dr. know what the output is at 7am every morning. This will allow your Dr. to know when to remove the drains.
2. The drain will typically be removed at the earliest when it is less than 10cc and is straw colored.
3. The drain should be kept on suction at all times. The way to do this is to squeeze the bulb with the white cap off, then screw the white cap on while squeezing. We will show you how to operate the drain before you leave after your procedure.

Food and Liquids

1. You should avoid hot (temperature) and spicy liquids and foods for the first 2 weeks. Drink plenty of fluids to make sure that your urine is not too dark. It should be a light yellow to clear in color. Eating helps you to absorb your medication and also prevent nausea with your pain medication. However, early on after surgery too much food can cause nausea

Medications

1. Follow instructions on prescribed medications
2. Never take Tylenol with Pain medicine at the same time.
3. Pain Medicine can cause constipation- drink plenty of fluids, Metamucil and or Colace as needed
4. **Always eat food with pain medicine and antibiotics**
5. No blood thinners, drinking alcohol, aspirin, anti inflammatories, high dose vitamin E, herbal medications or smoking for two weeks before & after the procedure.
6. In regards to pain medicine and your caretaker (if you are getting the procedure under local anesthesia you will not have a caretaker), the person taking care of you should stay up the whole night the first night or more as needed to make sure that you are breathing, to help you clean, taking medicines etc. You can take a half a pill of pain medication every hour until your pain gets to be 2 to 3 out of 10. You never want your pain to be zero out of 10 that means you're taking too much pain medication and you can potentially STOP BREATHING AND DIE. Also never take sleeping pills or other medications that make you tired while taking pain pills or you can potentially STOP BREATHING AND DIE. Also. you don't want your pain to be too high



AESTHETIC FACIAL BODY PLASTIC SURGERY

DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

(like 7-10) otherwise you'll have complications like bleeding, heart attacks, stroke, high blood pressure, death etc.

Activity

1.No strenuous activity, straining, bending over, lifting greater than 5 pounds for 2 weeks after surgery. It's okay to take a light walk for about 15-20 minutes once or twice a day after the first three days and increase your activity slowly after two weeks. **Light walks** also sometimes help with resolving residual bruising and swelling.

3. **You can bathe/shower after the first 72 hours**, just as long as you don't use very warm water. The water should be lukewarm just warm enough not to cause discomfort from being too cold.

5. Your healing, swelling and recovery will be 60% at 6 weeks and 80% at 6months. You should gauge your activity based on this. We generally advise starting regular activity slowly 2 weeks after your procedure

6. **AVOID THE SUN-** protect incision with SPF after incision is healed over. If you burn your incision, you most likely will have hyperpigmentation of the scar

7. Sleep with head elevated until swelling improves (2-3 pillows)

*Please text Dr Young daily closeup pictures for at least 2 weeks. Please send without vaseline nor betadine and after you've cleaned WITH FLASH. *

Patient

Date

Caregiver

Date



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

Post Procedure Instructions For Your Brow Lift And Wound Care (Long Form)

Congratulations on completing your Brow Lift procedure! Below is a list of instructions designed to help you through the recovery process. After you have read these instructions if you still have questions (especially if you think there is an emergency) please call 425-990-3223.

Please make sure to come in to see us the day after surgery for your post-op appointment. For your privacy we have a back entrance to be used for follow up appointments, there is a doorbell at this entrance. Please ring the bell upon your arrival and we will greet you at the door. We ask that you use this entrance for all follow up appointments for the first month after your procedure.

Things You Will Need:

- White Vinegar
- Distilled Water
- 3-4 Inch Ace Bandage (2 Better Than 1)
- Q-tips
- Vaseline
- 4 X 4 Inch Gauze
- Silicone Scar Healing Gel
- You can find links to specialized ice packs and facial | body garment wraps on our [Aesthetic Facial Body Plastic Surgery Patient Care Page](http://www.drphilipyoung.com/patient-care/) here:
<http://www.drphilipyoung.com/patient-care/>
- *If you can't afford these you can always use an ace bandage to wrap the areas and frozen peas for icing.*

Eye Care:

1. You should apply 1-2 drops of artificial tears into your eyes every hour to keep them moist.
2. Before you go to bed every night you should apply lacrilube into your eyes to lubricate your eyes and moisten them for the night while you sleep. This ointment



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

will cause blurriness and will eventually improve after a while when you wake up. This is essential to keeping your eyes moist after your procedure while you sleep. You can also use the lacrilube on your Blepharoplasty incisions instead of the Vaseline

General:

1. You should keep the original wrap that we put on until you come in the next day, at which time we will change it for you. You should then start cleaning your incisions at least 2-3 times a day or more to avoid scabbing and crusting.
2. After cleaning you should rewrap your face with the ace bandages and 4" x 4" gauze.
3. It is best to wrap your face for at least a week at all times (24 hours a day for 7 days).
4. After the first week, you should wrap your face at night for two months after your procedure.
5. After the first week you can start using the commercial wraps that can be purchased at the above mentioned sites.
6. For your drain: you should check the output of each drain separately, record and empty it every 6-8 hours. You should record how much the drainage is in cc or milliliter (ml) increments. The drain is optimally taken out when the output is less than 15-30cc(s) in a 24 hour period. The drain will typically be removed in the first 1-2 days after your procedure, but no later than 3-4 days after the procedure unless otherwise directed by your Doctor. The drain should be kept on suction at all times. The way to do this is to squeeze the bulb and then plug it to maintain the suction. If you have questions regarding this you can call us. We will show you how to operate the drain before you leave after the procedure. Please schedule an appointment to take out the drain no later than 3-4 days after your procedure.

Cleaning Your Incisions:

1. You should clean your incision 2-3 times a day or more for the first two weeks with 1 part vinegar and 3-4 parts distilled water mixture. This will help to avoid scabbing/crusting around the wound. You should clean more frequently if there is crusting. The goal is to have a very small | closely opposed incision line with no crusting for the best healing.
2. You should keep Vaseline on the incision at all times for the first 10-14 days or until further notified. This will allow your incision to heal faster. A light coat is suggested to help with cleaning.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

3. After two weeks you can consider Kelocote or other products containing silicone gel on the healed incisions. Apply the silicone gel for 12 hours a day on and 12 hours a day off for the next 3 months. The silicone gel only applies if you have an external incision. The other 12 hours that the silicone gel is off, you should apply spf 45 or greater sunscreen on your incisions (during the day)
4. More information on wound care can be found on this link:
<http://www.drphilipyoung.com/patient-care/procedure-process/wound-care-scar-prevention/>

Food and Liquids:

1. You should avoid hot (temperature) and spicy liquids and foods for the first 2 weeks. Drink plenty of fluids to make sure that your urine is not too dark. It should be a light yellow color. Eating helps you to absorb your medication and also prevent nausea with your pain medication. However, early on after surgery too much food can cause nausea.

Medications:

1. We recommend that you start taking your nausea and pain medications as soon as possible. Start with the nausea medication as soon as you can after surgery followed by the pain medication (1/2 tablet to one tablet) ½ hour later. You should only have a maximum of 12 tablets of Percocet (5/325) over 24 hours and no more. The pain medication takes an hour or so to take effect and during the start up time you will have pain. The nausea medication also takes about an hour to take effect.
2. Always try to remember not to take too much pain medication. Pain medications can slow your breathing, which can be dangerous. Also you can only have 4 grams of Tylenol in 24 hours; over this amount can lead to irreversible liver damage. So never combine regular Tylenol with your Pain medication.
3. Do not use high dose vitamin E, herbal medications or supplements, aspirin, Aleve, Ibuprofen or other anti-inflammatories for two weeks before and after your surgery. These medications and agents all cause bleeding.
4. You should start your antibiotics and Clonidine (if you have this medication) as soon as possible after your surgery as well. The antibiotics and Clonidine are to be taken regularly and finished completely. The nausea and pain medications are taken on an as needed basis.

Stitches:

1. Some of your stitches will need to be removed 6-7 days following your surgery; others may be removed a month later.
2. Sometimes stitches can break because we use very delicate stitches to get a good result. If this happens schedule an appointment as soon as possible for us to re-stitch the area.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

Bleeding:

1. You will have some mild bleeding from your incisions. This is normal when the local anesthesia wears off. The first 24 hours is the most common time period for severe bleeding. The other time point is at 7 days when the clots get dissolved. Hence the first two weeks are crucial and you should be careful.

Swelling:

1. We usually see you the next day after your procedure. This is when we determine if you have swelling that we should worry about.
2. Your swelling may increase over the next 3-4 days. Sometimes swelling lasts well over a week in a small population of patients. It all is dependent on the amount of surgery that you have.
3. Try to avoid sleeping on the sides of your face. This can lead to asymmetry and lead to a distorted look. Give yourself some time for the swelling to go down. All lumps and bumps eventually resolve.
4. Sleeping at a 30-45 degree angle, on your back can help the swelling for the first week.
5. Cool compresses (ice pack or bag of frozen peas) are essential for the first 7 days (or longer if needed) 15-20 minutes each hour. No warm compresses are recommended nor Massage unless directed by the Doctor.
6. 15-20 minute walks twice a day will also resolve residual bruising and swelling (start these 3 days after your surgery if you would like).
7. If you notice redness spreading, skin hot to the touch, white/yellow discharge from the incisions, or increased pain to one particular area, please set up an appointment to be seen this could be an infection that is easily treatable if caught early.

Sun Avoidance:

1. Sun avoidance is crucial to avoid having your wound turn dark from hyperpigmentation.
2. You must stay out of the sun or minimize your exposure for the next 6 months. If you do go out in the sun you must wear sunscreen SPF 45 and reapply every 30 minutes to an hour. You should try to avoid putting on sunscreen for the first two weeks.

Activity:

1. No strenuous activity, straining, bending over, lifting greater than 5 pounds, drinking alcohol or hot oral fluids, eating spicy foods, exercise, very hot showers and hot baths, aspirin, anti inflammatories, high dose vitamin E, herbal medications, smoking, for 2 weeks after surgery. It's okay to take a light walk for about 15-20



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

minutes once or twice a day after the first three days and increase your activity slowly after two weeks.

2. Light walks also sometimes help with resolving residual bruising and swelling.
3. You can bathe/shower after the first 72 hours, just as long as you don't use very warm water. The water should be lukewarm just warm enough not to cause discomfort from being too cold.
4. If you are straining going to the bathroom, in order to avoid pressure along your incision lines and to avoid bleeding, our Doctors usually suggest drinking a lot of fluids to help your bowel movements (to the point where your urine is clear), Metamucil daily, Colace 100mg orally twice a day. Our Doctors also suggest using suppositories or enema's as needed.
5. Your healing will be 60% at 6 weeks and 80% at 6 months. You should gauge your activity based on this. We generally advise starting regular activity slowly 2 weeks after your procedure.

Make Up:

1. Make up can usually be worn after 7 days depending on how your incisions heal. No makeup should be applied over raw areas.
2. Do not use age defying make-up around your incisions or on your eyes as it contains retinol and may cause an allergic reaction for up to one month.

Follow Up:

- i. We typically have most of our patients come back the next day.
- ii. You will then return in another 6-7 days after your surgery for suture removal unless otherwise directed.

If there any questions that are not being cleared up please call the office or set up an appointment immediately if you have concerns.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

PATIENT RIGHTS

The patient has the right to:

1. Treatment without regard to race, gender, age, national origin or cultural, economic, educational, or religious background, or the source of payment of his care.
2. Dignified, considerate and respectful care.
3. The knowledge of the name of the surgeon who has primary responsibility for coordination of his care and the names and professional relationships of other practitioners who will see him. All health care professionals practicing at the facility have had their credentials verified and have been approved to practice at the facility by the Governing Board.
4. Receives information from his surgeon about his illness, his course of treatment, and his prospects for recovery in terms that he can understand. When it is medically inadvisable to give such information to a patient, the information is provided to a person designated by the patient or to a legally authorized person.
5. Receive the necessary information about any proposed treatment or procedure in order to give informed consent or to refuse this course of treatment. Except in emergencies, this information shall include a description of all the procedure(s) or treatment(s), the medically significant risk(s) involved in the treatment, an alternate course of treatment or non-treatment, and the risk(s) involved in each, including the name of the person who would carry out the treatment(s) or procedure(s).
6. Participate actively in decision regarding his medical care. To the extent it is permitted by law, this includes the right to refuse treatment.
7. Full consideration of privacy concerning his medical care program. Case discussion, consultation, examination and treatment are confidential and shall be conducted discreetly. The patient has the right to be advised as to the reason for the presence of any individual.
8. Confidential treatment of all communications and records pertaining to his care. His written permission shall be obtained before his medical records are made available to anyone not concerned with his care.
9. Reasonable responses to any reasonable request he makes for services.
10. Reasonable continuity of care and to know, in advance the time and location of appointment(s), as well as, the practitioner providing the care.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

11. Be advised if the surgeon proposes to engage in or perform human experimentation affecting his care or treatment. The patient has the right to refuse to participate in such research projects.
12. Be informed by his surgeon, or designee, of his continuing health care requirements.
13. Examine and receive an explanation of his bill regardless of the source of payment.
14. Have all patients rights explained to the person who has legal responsibility to make decisions regarding medical care on behalf of the patient.
15. Express any grievances or suggestions verbally or in writing without fear of retribution or denial of care and expect his grievance to be reported to the person in charge immediately, and that his grievance will be investigated regarding treatment or care that is furnished, or fails to be furnished. The patient has the right to contact AAAHC.org.
16. Have information provided prior to 24 hours before the date of the procedure concerning the policies on advanced medical directives concerning such issues as living wills and durable powers of attorney that will be identified to the facility and followed as appropriate under State and Federal Regulations. Document in a prominent part of the patient's current medical record whether or not the individual has executed an advance directive.
17. Be informed of their right to change primary or specialty physicians if other qualified physicians are available.
18. Provided appropriate information regarding malpractice insurance coverage.
19. Patient will be treated respectfully regarding privacy, security, grievance resolution, spiritual care, and communication. If communication restrictions are necessary for patient care and safety, the facility must document and explain the restrictions to the patient and family.
20. Be protected from abuse and neglect.
21. Be informed of unanticipated outcomes.
22. Aesthetic Facial Body Plastic Surgery is owned and operated by Dr. Young.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

CONSENT FOR BROWLIFT

INSTRUCTIONS

This is an informed-consent document that has been prepared to help inform you of brow lift surgery, its risks, as well as alternative treatments.

It is important that you read this information carefully and completely. Please initial each page, indicating that you have read the page and sign the consent for surgery as proposed by your plastic surgeon and agreed upon by you.

GENERAL INFORMATION

The forehead and eyebrow region often show noticeable signs of aging. Looseness in these structures may cause drooping eyebrows, eyelid hooding, forehead furrows, and frown lines. In brow lift surgery, the structures responsible for these problems are tightened or altered to smooth the forehead, raise the upper eyebrows, and improve frown lines. A brow lift may be performed alone, or in conjunction with other procedures, such as a facelift or eyelid surgery.

Advances in brow lift surgery make it possible to perform the procedure through a variety of approaches, including endoscopy. Brow lift surgery is individualized for each patient. The surgical incisions used may vary with the technique selected by your surgeon to meet your needs. The brow lift cannot stop the process of aging.

ALTERNATIVE TREATMENTS

Alternative forms of treatment consist of not treating the laxness in the forehead and upper eyebrow region by brow lift surgery. Improvement of skin laxity and skin wrinkles may be attempted by other treatments or surgery such as laser resurfacing or chemical face peels. Risks and potential complications are associated with alternative surgical forms of treatment.

RISKS OF BROW LIFT SURGERY

Every surgical procedure involves a certain amount of risk and it is important that you understand these risks and the possible complications associated with them. In addition, every procedure has limitations. An individual's choice to undergo a surgical procedure is based on the comparison of the risk to potential benefit. Although the majority of patients do not experience these complications, you should discuss each of them with your plastic surgeon to make sure you understand all possible consequences of brow lift surgery.

Bleeding- It is possible, though unusual, to experience a bleeding episode during or after surgery. Should post-operative bleeding occur, it may require emergency treatment to drain accumulated blood (hematoma) or blood transfusion. Intra-operative blood transfusion may also be required. Do not take any aspirin or anti-inflammatory medications for ten days before or after surgery, as this may increase the risk of bleeding. Non-prescription "herbs" and dietary supplements can increase the risk of surgical bleeding. Hematoma can occur at any time following injury and may delay healing and cause scarring. If blood transfusions are necessary to treat blood loss, there is the risk of blood-related infections such as hepatitis and HIV (AIDS). Heparin medications that are used to prevent blood clots in veins can produce bleeding and decreased blood platelets.

Infection- Infection is unusual after surgery. Should an infection occur, additional treatment including antibiotics, hospitalization, or additional surgery may be necessary.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

Skin Discoloration / Swelling- Some bruising and swelling normally occurs following a brow lift. The skin in or near the surgical site can appear either lighter or darker than surrounding skin. Although uncommon, swelling and skin discoloration may persist for long periods of time and, in rare situations, may be permanent.

Change in Skin Sensation- It is common to experience diminished (or loss) of skin sensation in areas that have had surgery. Diminished (or loss) of skin sensation in the face and scalp area may not totally resolve after brow lift surgery. Chronic itching sensations can occur within the scalp and brow area following a brow lift.

Skin Contour Irregularities- Contour irregularities, depressions, and wrinkling of skin may occur after brow lift. Visible and palpable wrinkling of skin can occur. Residual skin irregularities at the ends of the incisions or "dog ears" are always a possibility and may require additional surgery. This may improve with time, or it can be surgically corrected.

Scarring- All surgery leaves scars, some more visible than others. Although good wound healing after a surgical procedure is expected, abnormal scars may occur within the skin and deeper tissues. Excessive scarring is uncommon. Scars may be unattractive and of different color than the surrounding skin tone. Scar appearance may also vary within the same scar. Scars may be asymmetrical (appear different on the right and left side of the body). There is the possibility of visible marks in the skin from sutures, staples, or hardware used during a brow lift. Additional treatments including surgery may be necessary to treat abnormal scarring.

Change in Surgical Approach for Brow Lift- In some situations, depending on factors discovered only at the time of surgery, your surgeon may have to make changes in surgical technique and approach to the brow lift procedure. This may require changing from an endoscopic (closed) technique to a standard (open) brow lift technique.

Surgical Anesthesia- Both local and general anesthesia involve risk. There is the possibility of complications, injury, and even death from all forms of surgical anesthesia or sedation.

Nerve Injury- There is the potential for injury to both motor and sensory nerves during a brow lift procedure. Weakness or loss in movements of the forehead or upper eyebrow may occur after surgery. Most individuals will notice a return of motor function; permanent weakness is rare. Injury may also occur in the sensory nerves of the forehead, scalp, and temple regions. Diminished sensation may normally occur in the scalp region after a brow lift surgery. Permanent numbness or painful nerve scarring is rare.

Damage to Deeper Structures- Deeper structures such as the eye, nerves, blood vessels, skull bone, muscles and lungs (pneumothorax) may be damaged during the course of surgery. The potential for this to occur varies according to the type of brow lift procedure being performed. Injury to deeper structures may be temporary or permanent.

Asymmetry- The human face is normally asymmetrical. There can be a variation from one side to the other in the results obtained from a brow lift procedure. Additional surgery may be necessary to attempt to revise asymmetry.

Delayed Healing- Wound disruption or delayed wound healing is possible. Some areas of the brow and scalp may not heal normally and may take a long time to heal. Some areas of skin may die, requiring frequent dressing changes or further surgery to remove the non-healed tissue. **Smokers have a greater risk of skin loss and wound healing complications.**



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

Allergic Reactions- In rare cases, local allergies to tape, suture materials and glues, blood products, topical preparations or injected agents have been reported. Serious systemic reactions including shock (anaphylaxis) may occur to drugs used during surgery and prescription medications. Allergic reactions may require additional treatment.

Seroma- Fluid accumulations infrequently occur in between the skin and the underlying tissues. Should this problem occur, it may require additional procedures for drainage of fluid.

Eye Irritation- Irritation or dryness in the eyes may occur after a brow lift or when the patient has eyelid surgery performed at the same time.

Pain- You will experience pain after your surgery. Pain of varying intensity and duration may occur and persist after brow lift surgery. Chronic pain may occur very infrequently from nerves becoming trapped in scar tissue.

Skin Sensitivity- Itching, tenderness, or exaggerated responses to hot or cold temperatures may occur after surgery. Usually this resolves during healing, but in rare situations it may be chronic.

Hair Loss- Hair loss may occur within the scalp or surgical incisions. The occurrence of this is not predictable. Hair loss may resolve slowly or in rare cases be permanent.

Hardware and Deeper Sutures- Some surgical techniques use small screws or permanent deep sutures to help suspend brow structures. Sutures may spontaneously poke through the skin, become visible or produce irritation that requires removal. In very unusual circumstances, a screw could penetrate through the skull. Intracranial injury is rare, but possible. If this occurs, additional treatment may be necessary. It may be necessary to remove hardware or deeper sutures at a later time.

Eyelid Disorders- Disorders that involve abnormal position of the upper eyelids (eyelid ptosis), loose eyelid skin, or abnormal laxness of the lower eyelid (ectropion) can coexist with sagging forehead and eyebrow structures. Brow lift surgery will not correct these disorders. Additional surgical procedures may be necessary.

Shock- In rare circumstances, your surgical procedure can cause severe trauma, particularly when multiple or extensive procedures are performed. Although serious complications are infrequent, infections or excessive fluid loss can lead to severe illness and even death. If surgical shock occurs, hospitalization and additional treatment would be necessary.

Unsatisfactory Result- Although good results are expected, there is no guarantee or warranty expressed or implied, on the results that may be obtained. You may be disappointed with the results of brow lift surgery. This would include risks such as asymmetry, unsatisfactory surgical scar location, unacceptable visible deformities at the ends of the incisions (dog ears), loss of facial movement, poor healing, wound disruption, and loss of sensation. It may be necessary to perform additional surgery to attempt to improve your results.

Cardiac and Pulmonary Complications- Surgery, especially longer procedures, may be associated with the formation of, or increase in, blood clots in the venous system. Pulmonary complications may occur secondarily to both blood clots (pulmonary emboli), fat deposits (fat emboli) or partial collapse of the lungs after general anesthesia. Pulmonary and fat emboli can be life-threatening or fatal in some circumstances. Air travel, inactivity and other conditions may increase the incidence of blood clots traveling to the lungs causing a major blood clot that may result in death. It is important to discuss with your physician any past history of blood clots or swollen legs that may contribute to this condition. Cardiac complications are a risk



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

with any surgery and anesthesia, even in patients without symptoms. If you experience shortness of breath, chest pains, or unusual heart beats, seek medical attention immediately. Should any of these complications occur, you may require hospitalization and additional treatment.

ADDITIONAL ADVISORIES

Long-Term Results- Subsequent alterations in forehead and upper eyebrow appearance may occur as the result of aging, weight loss or gain, sun exposure, pregnancy, menopause, or other circumstances not related to brow lift surgery. Brow lift surgery does not arrest the aging process or produce permanent tightening of the forehead. Future surgery or other treatments may be necessary to maintain the results of a brow lift procedure.

Skin Disorders / Skin Cancer- A brow lift is a surgical procedure for the tightening of skin and deeper structures of the face. Skin disorders and skin cancer may occur independently of a brow lift.

Use of Fibrin Sealants "Tissue Glue" Advisory-

Fibrin sealants which are produced from heat-treated human blood components to inactivate virus transmission are used to hold tissue layers together at surgery and to diminish post-operative bruising following surgery. This product has been carefully produced from screened donor blood plasma for hepatitis, syphilis, and human immunodeficiency virus (HIV). These products have been used safely for many years as sealants in cardiovascular and general surgery. This product is thought to be of help in diminishing surgical bleeding by adhering layers of tissue together.

Previous Surgical Scars- The presence of surgical scars from previous facial surgery may limit the amount of skin tightening that can be produced.

Female Patient Information- It is important to inform your plastic surgeon if you use birth control pills, estrogen replacement, or if you believe you may be pregnant. Many medications including antibiotics may neutralize the preventive effect of birth control pills, allowing for conception and pregnancy.

Intimate Relations After Surgery- Surgery involves coagulating of blood vessels and increased activity of any kind may open these vessels leading to a bleed, or hematoma. Activity that increases your pulse or heart rate may cause additional bruising, swelling, and the need for return to surgery and control bleeding. It is wise to refrain from sexual activity until your physician states it is safe.

Smoking, Second-Hand Smoke Exposure, Nicotine Products (Patch, Gum, Nasal Spray)-

Patients, who are currently smoking, use tobacco products, or nicotine products (patch, gum, or nasal spray) are at a greater risk for significant surgical complications of skin dying, delayed healing, and additional scarring. Individuals exposed to second-hand smoke are also at potential risk for similar complications attributable to nicotine exposure. Additionally, smoking may have a significant negative effect on anesthesia and recovery from anesthesia, with coughing and possibly increased bleeding. Individuals who are not exposed to tobacco smoke or nicotine-containing products have a significantly lower risk of this type of complication.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

It is important to refrain from smoking at least 2 weeks before and after surgery and until your physician states it is safe to return, if desired.

Mental Health Disorders and Elective Surgery- It is important that all patients seeking to undergo elective surgery have realistic expectations that focus on improvement rather than perfection. Complications or less than satisfactory results are sometimes unavoidable, may require additional surgery and often are stressful. Please openly discuss with your surgeon, prior to surgery, any history that you may have of significant emotional depression or mental health disorders. Although many individuals may benefit psychologically from the results of elective surgery, effects on mental health cannot be accurately predicted.

Medications- There are many adverse reactions that occur as the result of taking over-the-counter, herbal, and/or prescription medications. Be sure to check with your physician about any drug interactions that may exist with medications which you are already taking. If you have an adverse reaction, stop the drugs immediately and call your plastic surgeon for further instructions. If the reaction is severe, go immediately to the nearest emergency room. When taking the prescribed pain medications after surgery, realize that they can affect your thought process and coordination. Do not drive, do not operate complex equipment, do not make any important decisions, and do not drink any alcohol while taking these medications. Be sure to take your prescribed medication only as directed.

ADDITIONAL SURGERY NECESSARY

There are many variable conditions that may influence the long-term result of brow lift surgery. Secondary surgery may be necessary to obtain optimal results. Should complications occur, additional surgery or other treatments may be necessary. Even though risks and complications occur infrequently, the risks cited are particularly associated with brow lift surgery. Other complications and risks can occur but are even more uncommon. The practice of medicine and surgery is not an exact science. Although good results are expected, there is no guarantee or warranty expressed or implied, on the results that may be obtained. In some situations, it may not be possible to achieve optimal results with a single surgical procedure.

PATIENT COMPLIANCE

Follow all physician instructions carefully; this is essential for the success of your outcome. It is important that the surgical incisions are not subjected to excessive force, swelling, abrasion, or motion during the time of healing. Personal and vocational activity needs to be restricted. Protective dressings and drains should not be removed unless instructed by your plastic surgeon. Successful post-operative function depends on both surgery and subsequent care. Physical activity that increases your pulse or heart rate may cause bruising, swelling, fluid accumulation and the need for return to surgery. It is wise to refrain from intimate physical activities after surgery until your physician states it is safe. It is important that you participate in follow-up care, return for aftercare, and promote your recovery after surgery.

HEALTH INSURANCE

Most health insurance companies exclude coverage for cosmetic surgical operations such as brow lift surgery or any complications that might occur from surgery. Please carefully review your health insurance subscriber information pamphlet. **Most insurance plans exclude coverage for secondary or revisionary surgery.**

FINANCIAL RESPONSIBILITIES

The cost of surgery involves several charges for the services provided. The total includes fees charged by your surgeon, the cost of surgical supplies, anesthesia, laboratory tests, and possible outpatient hospital charges, depending on where the surgery is performed. Depending on whether the cost of surgery is covered by an insurance plan, you will be responsible for necessary co-payments, deductibles, and charges not covered. The fees charged for this procedure do not include any potential future costs for additional procedures that you elect to have or require in order to revise, optimize, or complete your outcome.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

Additional costs may occur should complications develop from the surgery. Secondary surgery or hospital day-surgery charges involved with revision surgery will also be your responsibility. **In signing the consent for this surgery/procedure, you acknowledge that you have been informed about its risk and consequences and accept responsibility for the clinical decisions that were made along with the financial costs of all future treatments.**

DISCLAIMER

Informed-consent documents are used to communicate information about the proposed surgical treatment of a disease or condition along with disclosure of risks and alternative forms of treatment(s), including no surgery. The informed-consent process attempts to define principles of risk disclosure that should generally meet the needs of most patients in most circumstances.

However, informed-consent documents should not be considered all inclusive in defining other methods of care and risks encountered. Your plastic surgeon may provide you with additional or different information which is based on all the facts in your particular case and the current state of medical knowledge.

Informed-consent documents are not intended to define or serve as the standard of medical care. Standards of medical care are determined on the basis of all of the facts involved in an individual case and are subject to change as scientific knowledge and technology advance and as practice patterns evolve.

It Is Important That You Read The Above Information Carefully And Have All Of Your Questions Answered Before Signing The Consent On The Next Page.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

CONSENT FOR SURGERY / PROCEDURE or TREATMENT

1. I hereby authorize Dr. Philip Young and such assistants as may be selected to perform the following procedure or treatment: Brow Lift Surgery (Forehead Lift)

I have received the following information sheet: Informed Consent - Brow Lift Surgery

2. I recognize that during the course of the operation and medical treatment or anesthesia, unforeseen conditions may necessitate different procedures than those above. I therefore authorize the above physician and assistants or designees to perform such other procedures that are in the exercise of his or her professional judgment necessary and desirable. The authority granted under this paragraph shall include all conditions that require treatment and are not known to my physician at the time the procedure is begun.
3. I consent to the administration of such anesthetics considered necessary or advisable. I understand that all forms of anesthesia involve risk and the possibility of complications, injury, and sometimes death.
4. I acknowledge that no guarantee or representation has been given by anyone as to the results that may be obtained.
5. For purposes of advancing medical education, I consent to the admittance of observers to the operating room.
6. I consent to the disposal of any tissue, medical devices or body parts which may be removed.
7. I consent to the utilization of blood products should they be deemed necessary by my surgeon and/or his/her appointees, and I am aware that there are potential significant risks to my health with their utilization.
8. I authorize the release of my Social Security number to appropriate agencies for legal reporting and medical-device registration, if applicable.
9. I understand that the surgeons' fees are separate from the anesthesia and hospital charges, and the fees are agreeable to me. If a secondary procedure is necessary, further expenditure will be required.
10. I realize that not having the operation is an option.
11. IT HAS BEEN EXPLAINED TO ME IN A WAY THAT I UNDERSTAND:
 - a. THE ABOVE TREATMENT OR PROCEDURE TO BE UNDERTAKEN
 - b. THERE MAY BE ALTERNATIVE PROCEDURES OR METHODS OF TREATMENT
 - c. THERE ARE RISKS TO THE PROCEDURE OR TREATMENT PROPOSED

I CONSENT TO THE TREATMENT OR PROCEDURE AND THE ABOVE LISTED ITEMS (1-12). I AM SATISFIED WITH THE EXPLANATION.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

CONSCIOUS SEDATION | GENERAL ANESTHESIA CONSENT FORM

INSTRUCTIONS

This is an informed consent document which has been prepared to help inform you about the anesthesia options available to you for your surgical procedure, their risks, as well as alternative treatment(s).

GENERAL INFORMATION

Washington State Law guarantees that you have both the right and the obligation to make decisions concerning your health care. Your physician can provide you with the necessary information and advice, but as a member of the health care team you must enter into the decision-making process. This form has been designed to acknowledge your acceptance of treatment recommended by your physician.

TYPES OF ANESTHESIA OFFERED:

1. Local Sedation

A local anesthetic agent such as Novocaine, Lidocaine | Xylocaine, or Marcaine all with or without epinephrine is introduced in to the tissue through injection and produces a numbness that allows surgery to be carried out with little to no discomfort while the patient is awake.

2. Oral Sedation

Local sedation is used in conjunction with Valium (sedative), Percocet (pain medication), and Phenergan (anti-nausea medication) which are ingested orally. You will experience little to no discomfort; be aware and conscious but fairly relaxed; and responsive to voice.

3. Conscious Sedation with Registered Nurse and your Physician

Versed or Fentanyl is administered through an intravenous line by our nurse. Other agents that are deemed necessary (with the patient's safety and comfort as the utmost priority), may be used including, but not limited to, ketamine, propofol, robinul, lebatolol, flumazenil, naloxone, ondansetron, valium, dexamethasone, etc. This level is a little deeper state of



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

sedation. You will be very sleepy but still responsive to more forceful amounts of voice and touch. Some refer to this type of sedation as Twilight.

4. Conscious Sedation with a Certified Registered Nurse Anesthetist

The last type of anesthesia offered is administered through an intravenous line given by a Certified Registered Nurse Anesthetist. This level can allow you to go into a deeper state of conscious sedation where you require more stimulation through touch and voice to become responsive. With this level of sedation, you will be much less aware of what is going on during the procedure than the other levels.

5. General Anesthesia with a Certified Registered Nurse Anesthetist or Anesthesiologist

This type of anesthesia offered is done through both an Intravenous line and an endotracheal tube (breathing tube down the windpipe). This level will take you to deepest level of sedation. You will not be aware of anything and will be in a deep sleep. We use the same medication as 3 and 4 options and also some anesthesia gases.

RISKS OF ANESTHESIA

I have been informed how each type of anesthesia is performed. I understand that all sedation and anesthesia medications involve risks of complications and serious possible damage to vital organs such as the brain, heart, lung, liver, and kidney, and that in some cases use of these medications may result in paralysis, cardiac arrest, and/or brain death from both known and unknown causes. I have been informed of possible alternative forms of treatment, including non-treatment.

I understand that, during the course of anesthesia, operation, post-operative care, medical treatment, or other procedures, unforeseen conditions may necessitate additional or different procedures than set forth above. I therefore authorize my above-named physician, and his/her assistants or designees, to perform such procedures that are considered necessary and desirable, in their professional judgment. The authority granted under this paragraph shall extend to the treatment of all conditions that require treatment and are not known to my physician at the time the medical or surgical procedure is commenced.

I consent to the administration of sedation or anesthesia by my attending physician, by an anesthesiologist, or other qualified party under the direction of a physician as may be deemed necessary.

I hereby authorize my physician / Aesthetic Facial Body Plastic Surgery and/or such associates or assistants as may be selected by said physician to administer anesthesia.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

I certify that my physician has informed me of the nature and character of the proposed treatment, of the anticipated results of the proposed treatment, of the possible alternative forms of treatment, and of any recognized serious possible risks and complications of the proposed treatment and of alternative forms of treatment, including non-treatment.

I CERTIFY THAT I HAVE HAD THE OPPORTUNITY TO ASK QUESTIONS, I HAVE HAD ALL ASPECTS OF THIS MEDICAL TREATMENT EXPLAINED TO MY SATISFACTION, AND I CONSENT TO THE USE OF CONSCIOUS SEDATION.

****Consent to be Signed Electronically as Part of the Medical Record****

Consent Signature - Patient Signs Here



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

CARETAKER CONSENT FORM

Your post-operative care is critical. Once you leave our office, your care will no longer be in our control and, therefore, you must have someone watching you carefully who can provide you the right doses of medicines. That is why our doctors recommend that you have 24 hour nursing care from a qualified nursing center. If instead you choose to have a family member or friend watch over you, then you must ensure that the person you select is qualified to take care of you during this critical state. Failure to have proper post-operative care may result in slowing your recovery, permanent damage and even death. Do not take the appointment of your caretaker lightly.

We reserve the right to send you to an aftercare facility if we deem that your caretaker is not of sufficient status to care for you in the first 24 hours.

FOR PATIENT TO SIGN

I _____ (full name) appoint _____ (full name) as my post-operative caretaker for my surgery on _____ (date). I understand that the doctors recommend that I have qualified 24 hour nursing care, but I choose this person as my caretaker and accept the risk of my decision. I also understand that my private medical information will be disclosed to my caretaker as needed to help with my recovery. I also understand that any failure on the part of my caretaker does not create a liability to AFBPS, which is not responsible for my choice in caretaker and his/her abilities. I remain solely responsible for my decision.

Patient Signature: _____ Date: _____

FOR CARETAKER TO SIGN

I _____ (full name) agree to care for _____ during the post-operative period of 24 hours or more as necessary after surgery on _____ (date). I do not take this obligation lightly and understand that I could be liable for failure to care for the patient properly. I will keep patient's medical information confidential and will not disclose said information to anyone except those people involved in patient's care. I agree to monitor the patient's vital signs by doing the following and keeping a record of my care:



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

Staying in the same room as the patient;

Making sure breathing is strong;

Asking questions to make sure patient is able to respond;

Making sure that the patient uses the restroom regularly;

Giving liquids and food as directed;

Giving the proper doses of medicine and recording patient's response.

All of these measures should be done on a regular basis over the course of the night. The intervals can be as frequent as every 5-15 minutes depending on the condition of the patient that you are caring for. If I have any question at all, I will call the doctor at 425 990 3223 or other numbers that are supplied to me. If there is any problem, I will immediately call 911 and the doctor at 425 990 3223 or other numbers that are supplied to me..

I also understand that signing this form does not create a relationship between myself and AFBPS; instead, my sole relationship is with the patient, who has chosen me to be the caretaker. Any failure on my part does not make AFBPS liable in any way.

My phone numbers are: _____

Caretaker signature: _____ Date: _____



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

PAIN MANAGEMENT AGREEMENT

I understand that I have a right to comprehensive pain management along with the surgery that I will undergo. I wish to enter into a treatment agreement to prevent possible chemical dependency. I understand that failure to follow any of these agreed statements might result in Aesthetic Facial Body Plastic Surgery, PLLC ("AFBPS") and their physicians to not provide ongoing care for me.

I agree to undergo pain management by Aesthetic Facial Body Plastic Surgery, PLLC. Pain Management provided by AFBPS is for the purpose of post-operative plastic surgery. I agree to the following statements:

I will not accept any narcotic prescriptions from another doctor unless approved by all physicians.

I will be responsible for making sure that I do not run out of my medications on weekends and holidays, because abrupt discontinuation of these medications will cause severe withdrawal syndrome.

I will only take the medication as directed by AFBPS.

I understand that I must keep my medications in a safe place.

I understand that AFBPS will not supply additional refills for the prescriptions of medications that I may lose.

If my medications are stolen, AFBPS will refill the prescription one time only if a copy of the police report of the theft is submitted to the physician's office.

I will not give my prescriptions to anyone else.

I will only use one pharmacy.

I will keep my scheduled appointments with AFBPS unless I give notice of cancellation 24 hours in advance.

I understand that pain medications can affect my breathing and could lead to life threatening situations if I am not careful. I understand that I should not take too much medication that I am too tired or drowsy that will lead me to stop breathing and result in death. I understand that I have a maximum amount that I can take in a 24 hour period but that some people respond differently and this maximum may be less than what is stated and that I need to see how the medication is affecting me.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

I understand that if there is any question or concern regarding taking pain medications or taking too much pain medication then I will contact your Doctor immediately or call 911.

I agree to refrain from all mind/mood altering/illicit/addicting drugs including alcohol unless authorized by AFBPS.

My treatment plan may change based on outcome of treatment, especially if pain medications are ineffective. Such medications will be discontinued.

I understand that AFBPS believes in the following "Pain Patients' Bill of Rights."

You have the right to:

Have your pain prevented or controlled adequately.

Have your pain and medication history taken.

Have your pain questions answered.

Know what medication, treatment or anesthesia will be given.

Know the risks, benefits and side effects of treatment.

Know what alternative pain treatments may be available.

Ask for changes in treatments if your pain persists.

Receive compassionate and sympathetic care.

Receive pain medication on a timely basis.

Refuse treatment without prejudice from your physician.

Include your family in decision-making.

Termination Clauses

The doctor may terminate this agreement at any time if he/she has cause to believe that I am not complying with the terms of this agreement, or to believe that I have made a misrepresentation or false statement concerning my pain or my compliance with the terms



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

of this agreement. Proof or verification of such beliefs is not required for termination and discontinuation of care.

I understand that I may terminate this agreement at any time.

If the agreement is terminated, I will not be a patient of AFBPS or your particular physician and would strongly consider treatment for chemical dependency if clinically indicated.

Patient/Guardian Signature:

Date

Print Patient/Guardian Name:

Date

Physician Signature: _____

Print Physician Name: _____



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

GENERAL, HIPAA, PHOTO | VIDEO INFORMED CONSENT FORM, AND RELEASE AGREEMENT

Aesthetic Facial Body Plastic Surgery, PLLC's ("AFBPS"), by and through Dr. Phillip Young, agree to

provide treatment to: _____ ("Patient" or "you") [insert Patient's name]

pursuant to terms and conditions set forth under this General Informed Consent Form and Release Agreement (the "Agreement") and such other consent or release AFBPS may require from time to time.

Patient has received materials, literature and documents regarding AFBPS's policies and guidelines for pre- and post-procedure activities and prohibitions, as well as medications to avoid and release of rights, including but not limited to the following:

1. Healing Body and Mind;
2. Your Anesthesia Experience;
3. Pre-Procedure Instructions;
4. Medications to Avoid;
5. Post-Procedure Instructions;
6. Post-Operative Instructions for Your Specific Procedure that you are receiving;
7. Patient Rights;
8. Anesthesia Consent Form;
9. Caretaker Consent Form;
10. Pain Management Consent Form; and
11. Photographic / Videographic Documentation Consent Form



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

By executing this Agreement, Patient certifies that he/she has: (i) read; (2) understood; and (3) had an opportunity to ask questions regarding each section of this Agreement and all materials, literature and documents provided by AFBPS. Patient understands that for each specific procedure, he/she will be required to sign additional consent forms addressing the specific risks, side effects, post-procedure care, etc., associated with those particular procedures Patient will undergo while under the care of AFBPS. If the person signing as the "Patient" under this Agreement is doing so on behalf of a minor, then such person certifies that he or she is the parent, guardian, or conservator of the minor and that such person is authorized to sign this consent form on the minor's behalf.

SECTION 1

INTRODUCTION TO AESTHETIC FACIAL BODY PLASTIC SURGERY, PLLC

Aesthetic Facial Body Plastic Surgery, PLLC is a Professional Service Corporation which performs various plastic surgery procedures to enhance facial aesthetics of its patients. These procedures can help to reduce the visible signs of aging, but cannot stop the process of aging. Since each individual's body is different, the risks and results of any medical procedure may vary from person to person. These procedures are generally performed under local, oral or conscious sedation and some individuals may need extra healing time and may not be able to return to work or normal activities for a prolonged period of time.

SECTION 2

ALTERNATIVES TO TREATMENT

There are surgical and nonsurgical methods for improving facial aesthetics and AFBPS will provide you with options and alternatives that may be suitable for your objectives, which you should carefully review with your treating physician before deciding on one or more treatment procedures.

SECTION 3

RISKS OF PROCEDURES

Every medical and surgical procedure involves a certain amount of risk and it is important that you understand these risks. An individual's choice to undergo a medical or surgical procedure is based on, among other things, the comparison of the risk to potential benefit. Although the majority of patients do not experience complications,



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

you should discuss each of them with your physician to make sure you understand the potential risks, complications, and consequences of the associated procedures. Whenever the skin is cut or punctured, it heals with a scar. Some procedures will result in a permanent scar.

Normal symptoms that occur during the recovery periods: swelling and bruising, discomfort and some pain, crusting along the incision lines, numbness of operated upon skin lasting 3 months or possibly longer or permanent, itching, redness of scars. With each individual procedure, the specific consent to perform the procedure will outline in more detail some of the symptoms, side effects and risks associated with such a procedure.

SECTION 4

POST-PROCEDURE CARE

Post-Procedure care is an important part of your plastic surgery experience. It is your obligation to make sure that you keep all your post-procedure appointments as directed and make sure that you promptly contact your physician and seek emergency care in case of a medical emergency. You must have a caretaker for the first 24 hours. **You should also record how you are taking your medications. You should record the date and time of each prescription drug you are taking, how much and what medications are given, and the total amounts of the drugs that are left each and every time.** Medications (especially pain medications) can be dangerous and you need to strictly follow the instructions on the prescription attached to the bottle.

SECTION 5

FINANCIAL POLICY REGARDING REVISION AND COMPLICATIONS

As you have been, or will be, advised, no plastic surgeon can guarantee a specific result. From time to time, some patients may require additional surgery to deal with revisions or complications. In cosmetic procedures, there are certain problems that are unavoidable regardless of quality of the care provided and diligence exercised by the doctor and his/her team.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

Examples of problems that a patient may encounter include bleeding and/or an unfavorable scar after a surgical procedure. In both of these cases, the patient may require additional surgery, either on an emergency basis (as in the case with bleeding) or an elective basis (as in the case of scarring).

We hope that no complication arises and no revisionary surgery becomes necessary in your case. However, no plastic surgeon can make such a guarantee to any of his or her patients. It is important for the patient undergoing an elective surgical procedure to understand that surgical revisions and complications may result in additional costs. Revisions within six (6) months from the original procedure date will not incur additional physician fee; but facility, anesthesia and other fees and costs shall be the sole responsibility of the patient. Notwithstanding the foregoing, any revisions after six (6) months of the original procedure date will incur all standard fees and costs.

If you have any questions regarding this policy, our office staff would be happy to discuss it with you.

SECTION 6

DEPOSIT | FEE | CANCELLATION

POLICY **Deposit | Fee for Surgical**

Procedures:

Procedure quotes are valid for 3 months. To ensure you receive the procedure pricing, your procedure must be scheduled within 3 months and completed within 12 months of receipt of the original quote. Based on years of experience and to ensure an efficient schedule for the health of our office, a non-refundable deposit of 10% of the total cost of the procedure is required to reserve your procedure date. **If you decide to reschedule or cancel your procedure you will be charged this deposit and it will be non-refundable and not applicable for future procedures.** To reschedule to a different date you will be required to place another non-refundable deposit of 10% of the total cost of the procedure.

The total cost of your procedure will be collected in full at the time of your Pre-Operative appointment 2 weeks before your procedure date. If you cancel or reschedule your procedure within the 2 week time frame you will be charged 50% of the total cost of your procedure. After your procedure, there are no refunds given. If you have any questions regarding our financial or refund policy, feel free to contact our Patient Care Coordinator or Office Manager. You may make a payment by contacting our office at 425.990.3223 or through our [Plastic Surgery Financing and Online Payment page](#).



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

Appointments | in-office procedures:

A deposit of \$250 / \$500 (respectively) will be collected at the time of scheduling a Botox or Filler appointment with Dr. Young. If you decide to reschedule or cancel your procedure you will be charged this deposit and it will be non-refundable and not applicable for future procedures. To reschedule to a different date you will be required to place another non-refundable deposit. For Botox, Dr Young has a 40 unit minimum and a 2 syringe minimum for Fillers.

The remaining cost of your treatment will be collected in full at the time of your appointment. After your treatment appointment, there are no refunds given. If you have any questions regarding our financial or refund policy, feel free to contact our Patient Care Coordinator or Office Manager. You may make a payment by contacting our office at 425.990.3223 or through our [Financing and Online Payment page](#). The deposit and other fees can be paid on our Financing and Online Payment page or by contacting our office at 425.990.3223.

SECTION 7

DISCLAIMERS, RELEASES AND COVENANTS

Computer imaging may be used during your consultation. Although we strive to achieve the very best results every time, these images are used to help guide us during your procedure and are not a guarantee of results.

You understand that AFBPS will request or require you to sign the following consent forms:



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

- Patient HIPAA Consent Form;
- General Instruction Form;
- Photographic/Videographic Documentation Consent Form;
- Pain Management Agreement;
- Caretaker Consent; and
- Consent forms for each individual procedure you will undergo while under the care of AFBPS.

Informed consent documents are used to communicate information about the proposed medical or surgical treatment along with disclosure of risks and alternative forms of treatment(s). The informed consent process attempts to define principles of risk disclosure that should generally meet the needs of most patients in most circumstances.

However, informed consent documents should not be considered all inclusive in defining other methods of care and risks encountered. Your physician may provide you with additional information, which is based on all the facts in your particular case and the state of medical knowledge.

Informed consent documents are not intended to define or serve as the standard of medical care. Standards of medical care are determined on the basis of all the facts involved in an individual case and are subject to change as science, knowledge, and technology advance and as practice patterns evolve.

For purposes of advancing medical education, you consent to the admittance of observers to the operating room.

You consent to the disposal of any tissue, medical device or body parts which may be removed.

You understand that the success of the procedure is to a great extent dependent upon your closely following Pre-Op and Post-Op instructions your doctor has provided to you. Post-Op care, activities and precautions have been explained to you and you understand them fully.



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

You also consent to the administration of such anesthetics as may be considered necessary and advisable by the attending physicians and/or anesthesiologist. You are aware that risks are involved with anesthesia, such as allergic or toxic reactions and even cardiac or respiratory arrest.

Your physician, and/or your physician's designees, reserve the right to discuss your case with any third parties if, in your physician's considered opinion, it becomes necessary to do so. Your signature below will indicate your consent to this reservation.

You have had sufficient opportunity to discuss your treatment with your physician and/or your physician's associates, and all your questions have been answered to your satisfaction. You believe that you have adequate knowledge upon which to give an informed consent to the proposed treatment.

SECTION 8
MOTOR VEHICLE AND PROCEDURE DATE POLICY

You are advised not to operate a motorized vehicle or power equipment on the day of surgery. The drugs administered during the procedure may impair driving ability and you should not drive when you are on any sedating medications such as sleeping pills, antihistamines, muscle relaxants, anti-anxiety medications, clonidine, and pain medications. AFBPS recommends that you have someone drive you to and from our facility the day of your procedure, if you are taking pain or sedation medications.

You hereby release and hold AFBPS and Dr. Phillip Young harmless from any and all actions, loss or injury sustained by you or any third party as a consequence of your operation of any motorized vehicle or equipment while under the influence of sedating medications prescribed to you.

SECTION 9
SMOKING

NO SMOKING FOR AT LEAST TWO (2) WEEKS BEFORE AND AFTER YOUR PROCEDURE!!! You have been informed by AFBPS that you are not to smoke for at least two (2) weeks before and after your scheduled procedure at AFBPS. If you are unable to maintain this nonsmoking policy before the procedure, then you must notify AFBPS immediately to reschedule your procedure date. If you are unable to maintain the nonsmoking policy after your procedure, then you must notify AFBPS and your doctor immediately to assess your health risk and seek appropriate medical attention as necessary. You understand that this policy is in place for your health and safety and you shall not hold AFBPS and Dr. Phillip Young responsible for any negative result which



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

may have been directly or indirectly caused by smoking.

You hereby attest that you have read and understood the above information carefully and have had all your questions answered before signing the consent form.

SECTION 10

ADVANCED MEDICAL DIRECTIVE

You acknowledge that you have been informed that your Advanced Medical Directive will be suspended while you are being treated at AFBPS. You have given a copy of your Advanced Medical Directive document to the staff at AFBPS; in the event that it is necessary that you be transferred to a hospital for acute care, every effort will be made to assure that a copy of this document will accompany you. You understand that it is not the responsibility of AFBPS to advise each care provider (emergency responders, emergency room, acute care facility, etc.) of your Advanced Medical Directive and that you should keep a copy of your Advanced Medical Directive with you and your designated health care proxy should also maintain a copy of the form.

If no copy of the Advanced Medical Directive is supplied for your medical record, you release AFBPS from any obligation or responsibility related to your status in this regard.

SECTION 11

CONSENT TO DRAW LABS FOR EXPOSURES

By signing this consent I also allow Aesthetic Facial Body Plastic Surgery and its Staff to carry out necessary blood work in the event of an accidental needle stick. The purpose of this is to allow Aesthetic Facial Body Plastic Surgery and its Staff to test your blood to see if you are a carrier of certain types of diseases including, but not limited to, Human Immunodeficiency Virus, Hepatitis, Syphilis, etc.

SECTION 12

Patient HIPPA Consent Form

Your health and health care information is both personal and private. Aesthetic Facial Body Plastic Surgery, P.S. is dedicated to protecting your health care information. This HIPPA Consent Form provides information about how Aesthetic Facial Body Plastic Surgery, P.S. may use and disclose your Protected Health Information (PHI).

As part of your medical treatment, Aesthetic Facial Body Plastic Surgery, P.S. originates and maintains



AESTHETIC FACIAL BODY PLASTIC SURGERY

DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

paper and/or electronic records which contain PHI such as: demographic information; personal and family histories; symptoms; examination and test results; diagnoses; past, present and future plans for care and treatment; and information received from other health care providers, your employer and any health care plan. Aesthetic Facial Body Plastic Surgery, P.S. maintains Privacy Practices and Policies regarding the disclosure of PHI.

The Patient understands that:

- Protected Healthcare Information may be disclosed or used for treatment, billing and payment, or healthcare operations;
- The patient has the right and the opportunity to review Aesthetic Facial Body Plastic Surgery, P.S.'s

Privacy Practices and Policies;

- Aesthetic Facial Body Plastic Surgery, P.S. reserves the right to change it's Privacy Practices and Policies at any time;
- The Patient has the right to request, in writing, restricted disclosure of their PHI, however,

Aesthetic Facial Body Plastic Surgery, P.S. is not bound by the restrictions unless an agreement regarding the requested restrictions has been reached;
- The Patient understand that they will be responsible for copying and mailing charges associated with sending their medical records.
- The patient may revoke their consent, in writing, at any time regarding all *future* disclosures.

By signing this form, you consent to our use and disclosure of protected health information about you for treatment, payment, or healthcare operations. You have a right to revoke this consent in writing, signed by you and delivered to our office. Revocation will apply to any future disclosures but not to any disclosure already made in reliance on your prior consent or as required by law. The practice provides this form to comply with the Health Insurance Portability and Accountability Act of 1996 (HIPAA). Aesthetic Facial Body Plastic Surgery, P.S. reserves the right to change its Privacy Practices and Policies at any time. A revised copy of the Privacy Practices and Policies may be requested by contacting the office.

SECTION 13

PHOTOGRAPHIC / VIDEOGRAPHIC DOCUMENTATION CONSENT FORM



AESTHETIC FACIAL BODY PLASTIC SURGERY

DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

I hereby give my consent to the taking of photographs and/or video by Aesthetic Facial Body Plastic Surgery, PLLC ("AFBPS") of me or parts of my body in connection with the procedure(s) to be performed by the physician at AFBPS for the sole purpose of internal use at AFBPS.

I provide this authorization as a voluntary, yet private contribution: (i) for use in my medical files - patient chart - at AFBPS; (ii) in the interests of the physician and office staff;

(iii) for the purpose of facilitating consultations and procedural explanations to/for me;

(iv) for AFBPS training purposes. I understand that such photographs shall become the property of AFBPS and may be retained by AFBPS but will not be released by AFBPS for any purposes such as print, visual or electronic media, medical journals and/or textbooks, or for the purpose of informing the medical profession or the general public about plastic surgery procedures and methods.

I understand that I may be asked to sign a separate consent in the future for the purpose of releasing my photos for other uses such as advertising for the rights of AFBPS, but will not be required to do so, and may refuse.

I understand that I may refuse to authorize the release of my photos for internal use and that my refusal to consent to the release will not affect the health care services I presently receive, or will receive, from AFBPS.

I understand that I have the right to inspect and copy the information that I have authorized to be disclosed. I further understand that I have the right to revoke this authorization in writing at any time, but if I do so it will not have any effect on any actions taken prior to my revocation.

I understand that the information disclosed, or some portion thereof, may be protected by state law and/or the federal Health Insurance Portability and Accountability Act of 1996 ("HIPAA").



AESTHETIC FACIAL BODY PLASTIC SURGERY
DR. PHILIP A. YOUNG, MD DR. RIKESH T. PARIKH, MD

I release and discharge AFBPS, the physicians, and all parties acting under the license and authority from all rights that I may have in the photographs and from any claim that I may have relating to such use in publication, including any claim for payment in connection with distribution or publications of the photographs.

I certify that I have read the above Authorization and Release and fully understand its terms. If signing on behalf of a minor, I certify that I am the parent, guardian, or conservator of the minor and I am authorized to sign this consent form on the minor's behalf.

****Consent Will Be Signed Electronically As Part of the Medical Record****